arXiv:1001.0058v1 [math.NT] 31 Dec 2009

A SUP-HODGE BOUND FOR EXPONENTIAL SUMS

CHUNLEI LIU

Abstract. The C-function of T-adic exponential sums is studeid. An explicit arithmetic bound is established for the Newton polygon of the C-function. This polygon lies above the Hodge polygon. It gives a sup-Hodge bound of the C-function of p-power order exponential sums.

1. Introduction

Let p be a prime number, $\mathbb{F}_p=\mathbb{Z}/(p)$, $\overline{\mathbb{F}}_p$ a fixed algebraic closure of $\mathbb{F}_p$, and $\mathbb{F}_{p^k}$ the subfield of $\overline{\mathbb{F}}_p$ with p^k elements.

Let $q>1$ be a power of p, W the ring scheme of Witt vectors, $\mathbb{Z}_q=W(\mathbb{F}_q)$, $\mathbb{Q}_q$ the fraction field of $\mathbb{Z}_q$, $\overline{\mathbb{Q}}_p=\lim\limits_{\overrightarrow{k}}\mathbb{Q}_{p^k}$, and $\mathbb{C}_p$ the p-adic completion of $\overline{\mathbb{Q}}_p$.

Let $\triangle\supsetneqq\{0\}$ be an integral convex polytope in $\mathbb{R}^n$, and I the set of vertices of $\triangle$ different from the origin. Let

$$f(x)=\sum_{u\in\triangle}(a_ux^u,0,0,\cdots)\in W(\mathbb{F}_q[x_1^{\pm1},\cdots,x_n^{\pm1}])\text{ with }\prod_{u\in I}a_u\neq0,$$

where $x^u=x_1^{u_1}\cdots x_n^{u_n}$ if $u=(u_1,\cdots,u_n)\in\mathbb{Z}^n$.

Definition 1.1. *For $k\in\mathbb{N}$, the sum*

$$S_f(k,T)=\sum_{x\in(\mathbb{F}_{q^k}^{\times})^n}(1+T)^{\mathrm{Tr}_{\mathbb{Z}_{q^k}/\mathbb{Z}_p}(f(x))}\in\mathbb{Z}_p[[T]]$$

is call a T-adic exponential sum. And the function

$$L_f(s,T)=\exp(\sum_{k=1}^{\infty}S_f(k,T)\frac{s^k}{k})\in1+s\mathbb{Z}_p[[T]][[s]],$$

as a power series in the single variable s with coefficients in the T-adic complete field $\mathbb{Q}_p((T))$, is called an L-function of T-adic exponential sums.

Let m be a positive integer, ζ_{p^m} a primitive p^m-th root of unity, and $\pi_m=\zeta_{p^m}-1$. Then $S_f(k,\pi_m)$ is the exponential sum studied by Liu-Wei [LWe]. If $m=1$, the exponential sum $S_f(k,\pi_m)$ was studied by Adolphson-Sperber [AS,AS2]. And, if $n=1$, the exponential sum $S_f(k,\pi_m)$ was studied by Kumar-Helleseth-Calderbank [KHC] and Li [Li].

This research is supported by NSFC Grant No. 10671015.

Definition 1.2. *The function*

$$C_f(s,T) = C_f(s,T;\mathbb{F}_q) = \exp(\sum_{k=1}^{\infty} -(q^k-1)^{-n} S_f(k,T)\frac{s^k}{k}),$$

as a power series in the single variable s with coefficients in the T-adic complete field $\mathbb{Q}_p((T))$, is called a C-function of T-adic exponential sums.

We have

$$L_f(s,T) = \prod_{i=0}^{n} C_f(q^i s,T)^{(-1)^{n-i+1}\binom{n}{i}},$$

and

$$C_f(s,T) = \prod_{j=0}^{\infty} L_f(q^j s,T)^{(-1)^{n-1}\binom{n+j-1}{j}}.$$

So the C-function $C_f(s,T)$ and the L-function $L_f(s,T)$ determine each other. From the last identity, one sees that

$$C_f(s,T) \in 1 + s\mathbb{Z}_p[[T]][[s]].$$

Let $C(\triangle)$ be the cone generated by $\triangle$, and $M(\triangle) = M(\triangle) \cap \mathbb{Z}^n$. There is a degree function deg on $C(\triangle)$ which is $\mathbb{R}_{\geq 0}$-linear and takes the values 1 on each co-dimension 1 face not containing 0. For $a \notin C(\triangle)$, we define $\deg(a) = +\infty$.

Definition 1.3. *A convex function on $[0,+\infty]$ which is linear between consecutive integers with initial value 0 is called the infinite Hodge polygon of $\triangle$ if its slopes between consecutive integers are the numbers $\deg(a)$, $a \in M(\triangle)$. We denote this polygon by $H_{\triangle}^{\infty}$.*

Liu-Wan [LWa] also proved the following.

Theorem 1.4 (Hodge bound)**.** *We have*

$$T-adic\ NP\ of\ C_f(s,T) \geq ord_p(q)(p-1)H_{\triangle}^{\infty},$$

where NP is the short for Newton polygon.

Denote by $\lceil x \rceil$ the least integer equal or greater than x, and by $\{x\}$ the fractional part of x.

Definition 1.5. *Let $C \subseteq M(\triangle)$ be a finite subset. We define*

$$r_C = \max_{\beta}(\#\{a \in C \mid \{\deg(pa)\}' \geq \beta\} - \#\{a \in C \mid \{\deg(a)\}' \geq \beta\}).$$

Definition 1.6. *Let $a \subseteq M(\triangle)$. We define*

$$\varpi(a) = \lceil p\deg(a)\rceil - \lceil \deg(a)\rceil + r_{\{a\in M(\triangle)|\deg(u)<\deg(a)\}\cup\{a\}} - r_{\{a\in M(\triangle)|\deg(u)<\deg(a)\}}.$$

Definition 1.7. *The arithmetic polygon $p_{\triangle}$ of $\triangle$ is a convex function on $[0,+\infty]$ which is linear between consecutive integers with initial value 0, and whose slopes between consecutive integers are the numbers $\varpi_{\triangle}(a)$, $a \in M(\triangle)$.*

One can prove the following.

Theorem 1.8. *We have then*

$$p_{\triangle} \geq (p-1)H^{\infty}_{\triangle}.$$

Moreover, they coincide at the point $n!\mathrm{Vol}(\triangle)$.

Let D be the least positive integer such that $\deg(M(\triangle)) \subseteq \frac{1}{D}\mathbb{Z}$. The main result of this paper is the following.

Theorem 1.9. *If* $p > 3D$, *then*

$$T-\text{adic NP of } C_f(s,T) \geq \mathrm{ord}_p(q)p_{\triangle}.$$

From the above theorem we shall deduce the following.

Theorem 1.10. *If* $p > 3D$, *then, for* $t \in \mathbb{C}_p$ *with* $0 \neq |t|_p < 1$, *we have*

$$t-\text{adic NP of } C_f(s,t) \geq \mathrm{ord}_p(q)p_{\triangle}.$$

2. The T-adic Dwork Theory

In this section we review the T-adic analogue of Dwork theory on exponential sums.

Let

$$E(t) = \exp(\sum_{i=0}^{\infty} \frac{t^{p^i}}{p^i}) = \sum_{i=0}^{+\infty} \lambda_i t^i \in 1 + t\mathbb{Z}_p[[t]]$$

be the p-adic Artin-Hasse exponential series. Define a new T-adic uniformizer π of $\mathbb{Q}_p((T))$ by the formula $E(\pi) = 1 + T$. Let $\pi^{1/D}$ be a fixed D-th root of π. Let

$$L = \{ \sum_{u \in M(\triangle)} c_u \pi^{\deg(u)} x^u : \ c_u \in \mathbb{Z}_q[[\pi^{1/D}]]\}.$$

Let $a \mapsto \hat{a}$ be the Teichmüller lifting. One can show that the series

$$E_f(x) := \prod_{a_u \neq 0} E(\pi \hat{a}_u x^u) \in L.$$

Note that the Galois group of $\mathbb{Q}_q$ over $\mathbb{Q}_p$ can act on L but keeping $\pi^{1/D}$ as well as the variable x fixed. Let σ be the Frobenius element in the Galois group such that $\sigma(\zeta) = \zeta^p$ if ζ is a $(q-1)$-th root of unity. Let Ψ_p be the operator on L defined by the formula

$$\Psi_p(\sum_{i \in M(\triangle)} c_i x^i) = \sum_{i \in M(\triangle)} c_{pi} x^i.$$

Then $\Psi := \sigma^{-1} \circ \Psi_p \circ E_f$ acts on the T-adic Banach module

$$B = \{ \sum_{u \in M(\triangle)} c_i \pi^{\deg(u)} x^u \in L, \ \mathrm{ord}_T(c_u) \to +\infty \text{ if } \deg(u) \to +\infty\}.$$

We call it Dwork's T-adic semi-linear operator because it is semi-linear over $\mathbb{Z}_q[[\pi^{\frac{1}{D}}]]$.

Let $b=\log_p q$. Then the b-iterate Ψ^b is linear over $\mathbb{Z}_q[[\pi^{1/D}]]$, since

$$\Psi^b=\Psi_p^b\circ\prod_{i=0}^{b-1}E_f^{\sigma^i}(x^{p^i}).$$

One can show that Ψ is completely continuous in the sense of Serre [Se]. So $\det(1-\Psi^b s\mid B/\mathbb{Z}_q[[\pi^{\frac{1}{D}}]])$ and $\det(1-\Psi s\mid B/\mathbb{Z}_p[[\pi^{\frac{1}{D}}]])$ are well-defined.

We now state the T-adic Dwork trace formula [LWa].

Theorem 2.1. *We have*

$$C_f(s,T)=\det(1-\Psi^b s\mid B/\mathbb{Z}_q[[\pi^{\frac{1}{D}}]]).$$

Lemma 2.2. *The Newton polygon of* $\det(1-\Psi^b s^b\mid B/\mathbb{Z}_q[[\pi^{\frac{1}{D}}]])$ *coincides with that of* $\det(1-\Psi s\mid B/\mathbb{Z}_p[[\pi^{\frac{1}{D}}]])$.

Proof. Note that

$$\det(1-\Psi^b s\mid B/\mathbb{Z}_p[[\pi^{\frac{1}{D}}]])=\mathrm{Norm}(\det(1-\Psi^b s\mid B/\mathbb{Z}_q[[\pi^{\frac{1}{D}}]])),$$

where Norm is the norm map from $\mathbb{Z}_q[[\pi^{\frac{1}{D}}]]$ to $\mathbb{Z}_p[[\pi^{\frac{1}{D}}]]$. The lemma now follows from the equality

$$\prod_{\zeta^b=1}\det(1-\Psi\zeta s\mid B/\mathbb{Z}_p[[\pi^{\frac{1}{D}}]])=\det(1-\Psi^b s^b\mid B/\mathbb{Z}_p[[\pi^{\frac{1}{D}}]]).$$

□

Write

$$\det(1-\Psi s\mid B/\mathbb{Z}_p[[\pi^{\frac{1}{D}}]])=\sum_{i=0}^{+\infty}(-1)^i c_i s^i.$$

Theorem 2.3. *The T-adic Newton polygon of* $\det(1-\Psi^b s\mid B/\mathbb{Z}_q[[\pi^{\frac{1}{D}}]])$ *is the lower convex closure of the points*

$$(m, ord_T(c_{bm})),\ m=0,1,\cdots.$$

Proof. By Lemma 2.2, the T-adic Newton polygon of $\det(1-\Psi^b s^b\mid B/\mathbb{Z}_q[[\pi^{\frac{1}{D}}]])$ is the lower convex closure of the points

$$(i,\mathrm{ord}_T(c_i)),\ i=0,1,\cdots.$$

It is clear that $(i,\mathrm{ord}_T(c_i))$ is not a vertex of that polygon if $b\nmid i$. So that Newton polygon is the lower convex closure of the points

$$(bm,\mathrm{ord}_T(c_{bm})),\ m=0,1,\cdots.$$

It follows that the T-adic Newton polygon of $\det(1-\Psi^b s\mid B/\mathbb{Z}_q[[\pi^{\frac{1}{D}}]])$ is the lower convex closure of the points

$$(m,\mathrm{ord}_T(c_{bm})),\ m=0,1,\cdots.$$

□

3. The arithmetic bound

In this section we prove the following.

Theorem 3.1. *We have*

$$\mathrm{ord}_T(c_{bm}) \geq p_{\triangle}(m).$$

Proof. First, we choose a basis of $B \otimes_{\mathbb{Z}_p} \mathbb{Q}_p(\pi^{1/D})$ over $\mathbb{Q}_p(\pi^{1/D})$ as follows. Fix a normal basis $\bar{\xi}_i$, $i \in \mathbb{Z}/(b)$ of $\mathbb{F}_q$ over $\mathbb{F}_p$. Let ξ_i be their Terchmüller lift of $\bar{\xi}_i$. The system ξ_i, $i \in \mathbb{Z}/(b)$ is a normal basis of $\mathbb{Q}_q$ over $\mathbb{Q}_p$. Then $\{\xi_i x^u\}_{u \in M(\triangle), 1 \leq i \leq b}$ is a basis of $B \otimes_{\mathbb{Z}_p} \mathbb{Q}_p(\pi^{1/D})$ over $\mathbb{Q}_p(\pi^{1/D})$.

Secondly, we write out the matrix of Ψ on $B \otimes_{\mathbb{Z}_p} \mathbb{Q}_p(\pi^{1/D})$ with respect to the basis $\{\xi_i x^u\}_{u \in M(\triangle), 1 \leq i \leq b}$. Write

$$E_f(x) = \sum_{u \in M(\triangle)} \gamma_u x^u,$$

and

$$\sigma^{-1}(\xi_j \gamma_{pu-w}) = \sum_{i=1}^{b} \gamma_{(u,i),(w,j)} \xi_i.$$

Then

$$\begin{aligned}\Psi(\xi_j x^w) &= \sum_{u \in M(\triangle)} \sigma^{-1}(\xi_j \gamma_u) \Psi_p(x^{u+w}) \\ &= \sum_{u \in M(\triangle)} \sigma^{-1}(\xi_j \gamma_{pu-w}) x^u \\ &= \sum_{u \in M(\triangle)} \sum_{i=1}^{b} \gamma_{(u,i),(w,j)} \xi_i x^u.\end{aligned}$$

So $(\gamma_{(u,i),(w,j)})_{u,w \in M(\triangle), 1 \leq i,j \leq b}$ is the matrix of Ψ on $B \otimes_{\mathbb{Z}_p} \mathbb{Q}_p(\pi^{1/D})$ with respect to the basis $\{\xi_i x^u\}_{u \in M(\triangle), 1 \leq i \leq b}$.

Thirdly, we claim that

$$\mathrm{ord}_T(\gamma_{(u,i),(w,j)}) \geq \lceil \deg(pu - w) \rceil.$$

In fact, this follows from the equality

$$\sigma^{-1}(\xi_j \gamma_{pu-w}) = \sum_{i=1}^{b} \gamma_{(u,i),(w,j)} \xi_i,$$

and the inequality $\mathrm{ord}_T(\gamma_u) \geq \lceil \deg(u) \rceil$.

Finally, we show that

$$\mathrm{ord}_T(c_{bm}) \geq p_{\triangle}(m).$$

Note that

$$c_{bm} = \sum_{A} \det((\gamma_{(u,i),(w,j)})_{(u,i),(w,j) \in A}),$$

where A runs over all subsets of $M(\triangle)\times\mathbb{Z}/(b)$ with cardinality bm. So it suffices to show that

$$\mathrm{ord}_T(\det(\gamma_{(i,u),(j,\omega)})_{(i,u),(j,\omega)\in A})\geq bp_{\triangle}(m).$$

Note that

$$\det(\gamma_{(i,u),(j,\omega)})_{(i,u),(j,\omega)\in A})=\sum_{\tau\in S_A}\sum_{a\in A}\mathrm{ord}_{\pi}(\gamma_{a,\tau(a)}),$$

where S_A is the permutation group of A. So it suffices to show that

$$\sum_{a\in A}\mathrm{ord}_{\pi}(\gamma_{a,\tau(a)})\geq bp_{\triangle}(m),\ \ \tau\in S_A.$$

Since

$$\mathrm{ord}_T(\gamma_{(u,i),(w,j)})\geq\lceil\deg(pu-w)\rceil,$$

the theorem follows from the following. ☐

Theorem 3.2. *If $p>3D$, A is a subset of $M(\triangle)\times\mathbb{Z}/(b)$ with cardinality bm, and $\tau\in S_A$, then*

$$\sum_{a\in A}\lceil\deg(p\nu(a)-\nu(\tau(a)))\rceil\geq bp_{\triangle}(m),$$

where $\nu(u,i)=u$.

Proof. We have

$$\sum_{a\in A}\lceil\deg(p\nu(a)-\nu(\tau(a)))\rceil\geq\sum_{a\in A}\lceil\deg(p\nu(a))-\nu(\tau(a))\rceil$$

$$\geq\sum_{a\in A}\lceil\deg(p\nu(a))\rceil-\lceil\deg(\nu(\tau(a)))\rceil+1_{\{\deg(p\nu(a))\}'>\{\deg(\nu(\tau(a)))\}'}$$

$$\geq\sum_{a\in A}\lceil\deg(p\nu(a))\rceil-\lceil\deg(\nu(a))\rceil+1_{\{\deg(p\nu(a))\}'>\{\deg(\nu(\tau(a)))\}'}.$$

Choose a set B of cardinality $|A|$ such that $B\cap A$ is as big as possible under the condition that, for some α,

$$\{a\in M(\triangle)\times\mathbb{Z}/(b)\mid\deg(\nu(a))<\alpha\}\subseteq B\subsetneq\{a\in M(\triangle)\times\mathbb{Z}/(b)\mid\deg(\nu(a))\leq\alpha\}.$$

Choose a permutation τ_0 on $B\cap A$ which agrees with τ on $(B\cap A)\cap\tau^{-1}(B\cap A)$. Extend it trivially to B. We have

$$\sum_{a\in A}\lceil\deg(p\nu(a))\rceil-\lceil\deg(\nu(a))\rceil\geq\sum_{a\in B}(\lceil\deg(p\nu(a))\rceil-\lceil\deg(\nu(a))\rceil)+2\#(A\backslash B).$$

We also have

$$\sum_{a\in A}1_{\{\deg(p\nu(a))\}'>\{\deg(\nu(\tau(a)))\}'}\geq\sum_{a\in B}1_{\{\deg(p\nu(a))\}'>\{\deg(\nu(\tau_0(a)))\}'}-2\#(A\setminus B).$$

It follows that

$$\sum_{a\in A}\lceil\deg(p\nu(a))\rceil-\lceil\deg(\nu(a))\rceil+1_{\{\deg(p\nu(a))\}'>\{\deg(\nu(\tau(a)))\}'}$$

$$\geq \sum_{a\in B}\lceil \deg(p\nu(a))\rceil - \lceil \deg(\nu(a))\rceil + 1_{\{\deg(p\nu(a))\}'>\{\deg(\nu(\tau_0(a)))\}'}$$

$$\geq \sum_{a\in B}(\lceil \deg(p\nu(a))\rceil - \lceil \deg(\nu(a))\rceil) + r_B,$$

where

$$r_B = \max_\beta(\#\{a\in B \mid \{\deg(p\nu(a))\}' \geq \beta\} - \#\{a\in B \mid \{\deg(\nu(a))\}' \geq \beta\}).$$

Choose a set C of cardinality m such that for some α,

$$\{a\in M(\triangle) \mid \deg(\nu(a)) < \alpha\} \subseteq C \subsetneq \{a\in M(\triangle) \mid \deg(\nu(a)) \leq \alpha\}.$$

Recall that

$$r_C = \max_\beta(\#\{a\in C \mid \{\deg(pa)\}' \geq \beta\} - \#\{a\in C \mid \{\deg(a)\}' \geq \beta\}).$$

It is easy to see that $r_B = br_C$, and

$$\sum_{a\in B}(\lceil \deg(p\nu(a))\rceil - \lceil \deg(\nu(a))\rceil) + r_B$$

$$= b\sum_{a\in C}(\lceil \deg(pa)\rceil - \lceil \deg(a)\rceil) + br_C = bp_\triangle(m).$$

The theorem now follows. □

References

[AS] A. Adolphson and S. Sperber, Newton polyhedra and the degree of the L-function associated to an exponential sum, Invent. Math. 88(1987), 555-569.

[AS2] A. Adolphson and S. Sperber, Exponential sums and Newton polyhedra: cohomology and estimates, Ann. Math., 130 (1989), 367-406.

[BF] R. Blache and E. Férard, Newton straitification for polynomials: the open stratum, J. Number Theory, 123(2007), 456-472.

[KHC] P. V. Kumar, T. Helleseth and A. R. Calderbank, An upper bound for some exponential sums over Galois rings with applications, IEEE Trans. Inform. Theory 41 (1995), no.2, 456-468.

[Li] W.-C. W. Li, Character sums over p-adic fields, J. Number Theory 74 (1999), no.2, 181-229.

[LWa] C. Liu and D. Wan, T-adic exponential sums, Algebra & Number theory, Vol. 3, No. 5 (2009), 489-509.

[LWe] C. Liu and D. Wei, The L-functions of Witt coverings, Math. Z., 255 (2007), 95-115.

[Se] J-P. Serre, Endomorphismes complétement continus des espaces de Banach p-adiques, Publ. Math., IHES., 12(1962), 69-85.

[Zh1] J. H. Zhu, p-adic variation of L functions of one variable exponential sums, I. Amer. J. Math., 125 (2003), 669-690.

[Zh2] J. H. Zhu, Asymptotic variation of L functions of one-variable exponential sums, J. Reine Angew. Math., 572 (2004), 219–233. 1529–1550.

Department of Mathematics, Shanghai Jiao Tong University, Shanghai 200240, P.R. China.

E-mail address: `clliu@sjtu.edu.cn`